

Bupivacaine: A Review

Charles R. Babst, D.D.S.*

Bert N. Gilling, M.D.**

Queen's Medical Center, Honolulu, Hawaii

ABSTRACT

A review of current significant literature concerning bupivacaine hydrochloride (Marcaine) is presented with particular emphasis on clinical use in oral surgery. The major advantages compared with other presently used local anesthetics are an increased duration of action and a favorable potency to toxicity ratio.

Bupivacaine HCL (1-butyl-2', 6' pipecoloxylidide hydrochloride)* is a long acting amide local anesthetic (Fig. 1). First synthesized in 1957 by Ekenstam at A. B. Bafors Laboratories in Mölndel, Sweden, this drug has undergone trials and varying degrees of acceptance in virtually every area of local anesthetic practice.¹




Figure 1
Chemical Structure of Bupivacaine

Mechanism of Action

The mechanism of action of bupivacaine is presumed to be the same as for other local anesthetics. Current local anesthetic theory holds that these compounds obstruct the inward flow of sodium ions through the nerve membrane, thus preventing the generation of an action potential.² Competitive binding to calcium sites is postulated to occur in the external lipid layer of the nerve membrane with resultant secondary interference of mobile phosphate groups. Passage of sodium ions is blocked by preventing molecular membrane reconfiguration from the resting (sodium impermeable) to the active (sodium permeable) state.

The increased duration of action of bupivacaine is ascribed to its affinity for nerve tissue.³

Clinical Use

Bupivacaine is utilized for intraoperative local anesthesia, post operative analgesia and in the treatment of chronic pain.

Bupivacaine is widely used in obstetrics. In lumbar epidural anesthesia the drug appears innocuous to mother and fetus.^{4,5} The acceptable therapeutic index is largely due to the small amount of drug needed per unit time. The indications for bupivacaine in obstetrical analgesia are enhanced by the insignificant motor blockade in concentrations less than 0.5%.^{6,7}

Caudal blocks with bupivacaine for vaginal delivery are more efficacious due to the increased duration of analgesia,^{8,9,10} however fetal deaths have been reported secondary to paracervical block.¹¹ This latter method of administration is contraindicated unless epidural block is incomplete or unavailable at a given institution.

Excellent sensory anesthesia is reported with 0.5% bupivacaine epidural blocks for thoracic and abdominal surgery.^{12,13} The increased duration of action postpones the patients initial request for post operative analgesics.¹³ Continuous thoracic epidural infusion of 1.0% bupivacaine does not provide analgesia of greater duration than lesser concentrations of bupivacaine.¹⁴ Furthermore, tachyphalaxis, extensive segmental spread, urinary retention, high plasma drug concentrations, and inadequate operative anesthesia may occur.

Intraoperative anesthesia by intercostal injection of bupivacaine is effective,^{15,16,17} with a duration of four¹⁵ to five¹⁶ hours.

Bupivacaine combined with 2-chlorprocaine for brachial plexus blocks provides rapid onset and an increased duration of anesthesia over 2-chlorprocaine alone.^{18,19} No toxic reactions and a quiescent post operative period are reported.^{18,19}

In ulnar nerve block, bupivacaine 0.25% or 0.5% provides excellent sensory and sympathetic anesthesia, but motor paralysis is not complete. Initial onset for sensory fibers is within eight minutes.²⁰

For intravenous regional anesthesia, bupivacaine provides rapid onset (3-5 minutes), good muscle relaxation, and fewer toxic reactions relative to lidocaine.²¹

*Dental Resident Queen's Medical Center, Honolulu, Hawaii.

**Assistant Clinical Professor of Surgery (Anesthesiology), University of Hawaii, School of Medicine.

*Available as Marcaine HCL, Winthrop Laboratories, New York, N.Y.

Bupivacaine has also been used in non-surgical procedures. Epidural administration has provided total resolution of non-malignant chronic pain.²³ In a limited study,²⁴ intra-gasserian ganglionic injection of bupivacaine moderated exacerbations of trigeminal neuralgia.

Clinical Effectiveness

Effectiveness of bupivacaine is evaluated by toxicity to potency ratio, latency of onset, degree of sympathetic, sensory, and motor block, duration of analgesia, and regression time. Statistical data and clinical results are inseparably related to the vascularity of the area of injection and the concentration and quantity of the agents used.^{25,26}

Bupivacaine 0.5% initially appeared approximately equitoxic-equipotent to 0.5% tetracaine,^{26,27} 2.0% lidocaine,^{5,26,28,29} 3.0% mepivacaine,^{26,27,28} and 1.0% etidocaine.^{6,7,16,29,30,31,32} More recent literature reveals a toxicity to potency ratio even more favorable than previously stated.³³ The onset of action of bupivacaine (2-10 minutes, variable with source and nature of injection) is similar to tetracaine²⁶ and mepivacaine²⁶ but is somewhat slower than lidocaine,^{26,34,35} prilocaine,²⁶ and etidocaine.^{7,20,36}

Sympathetic fibers are more easily blocked than sensory or motor fibers.²⁶ Sympathetic blocks may manifest themselves in clinical phenomena as Horner's syndrome³⁸ or hypotension.³⁷ Bupivacaine blocks sympathetic fibers more effectively than etidocaine.²⁰

Löfström²⁶ contends that in equitoxic-equipotent doses, bupivacaine provides as complete and profound a sensory block as tetracaine, lidocaine, prilocaine, and mepivacaine.

Bupivacaine, prilocaine, and mepivacaine all provide complete motor blocks.²⁶ Both tetracaine²⁶ and etidocaine^{6,7,20,31,32,36,39,40} produce a more profound motor block than bupivacaine.

Bupivacaine, tetracaine and etidocaine have much longer durations of action than other local anesthetics. In equitoxic-equipotent doses, bupivacaine shows longer and more profound sensory blocks than either tetracaine²⁶ or etidocaine.^{6,7,31,32,36,39} Peripheral nerve blocks^{26,35} of eight to twelve hours and intercostal blocks²⁶ or 14 hours are reported. Vasoconstrictors with bupivacaine do not increase the duration of anesthetic action.^{6,26,40,41}

Regression time (the time from reappearance of pain perception until complete recovery of pain sensation) is longer with bupivacaine than with other agents.^{26,35}

Tachyphylaxis appears less likely with this agent than with other local anesthetics because fewer injections are usually made.^{5,26}

Bioavailability

The biologic fate of bupivacaine is dependent on absorption, distribution, and elimination.⁴²

Absorption and distribution are influenced by the vascularity of the injection site, mode of injection, and the degree of drug ionization.⁴²

Highly vascularized sites contribute to an increased drug plasma concentration.^{25,42}

Intravenous administration provides a peak plasma concentration shortly after injection.⁴² At this point, the drug's half life is 45 minutes which corresponds to tissue redistribution. Four hours after IV administration, $T_{1/2}$ becomes $2\frac{1}{2}$ hours due to elimination. IV drip infusion results in a slower increase in plasma concentration and a tendency to plateau which is indicative of equilibrium.⁴² Injection for nerve block results in maximum plasma levels in 15 to 30 minutes.⁴³

Local anesthetics are weak bases and diffuse through membranes as a lipid soluble unionized moiety.⁴² Therefore, physiologic alterations such as acidosis, which change bupivacaine to its ionized form, will inhibit membrane transport, delay onset, decrease effectiveness, decrease plasma concentrations, decrease toxicity, and decrease elimination via decreased delivery to liver and kidneys.⁴²

As with other amide local anesthetics⁴⁴ bupivacaine is mainly metabolized in the liver by N-dealkylation and glucuronide conjugation of the hydroxylated parent compound.⁴⁵ Time for metabolism of bupivacaine is equal to lidocaine and etidocaine.⁴⁶ Elimination is mainly via urinary excretion; however, some excretion of metabolites via the lungs and bile undoubtedly occurs.⁴⁷ Very little unchanged drug is recovered in the urine.^{45,48}

Bupivacaine is highly bound to non-albumin plasma proteins.⁴² This binding is more extensive than either lidocaine or mepivacaine,⁴² but equal to etidocaine.⁴⁹ Subsequent distribution of bupivacaine and etidocaine is greater than lidocaine and mepivacaine.⁴⁶ This increased protein binding results in a decreased tissue to blood coefficient. Therefore, the drug is presented to the liver quickly, but hepatic uptake is slow.⁴² Protein binding contributes to a decreased fetal to maternal drug concentration ratio, which increases the safety of this agent in obstetrics.¹⁰ Epinephrine decreased fetal bupivacaine levels further in one study.⁵⁰

The bound fraction of bupivacaine is pharmacologically inactive; it is the unbound fraction which is responsible for toxic reactions.⁴² Demerol, diphenylhydantoin, quinidine, and desipramine all readily displace bupivacaine from protein binding sites.⁵¹ However, the apparent increased potential for bupivacaine toxicity is not realized since erythrocyte binding sites readily absorb the displaced drug.⁵¹

Toxicity

Systemic local toxicity manifests itself in slurred speech, nystagmus, twitching, disorientation, circumoral numbness, lightheadedness, paresthesias, drowsiness, and convulsions.^{38,52} These CNS effects usually manifest themselves before ventilatory or circulatory compromise.^{53,54}

Toxicity increases with increasing rate of epidural infusion.^{29,52} Intermittant epidural block increases effectiveness and decreases toxicity compared with continuous infusion.³⁸ Bupivacaine appears slightly more

toxic than etidocaine which is slightly more toxic than lidocaine. Despite the differences in absolute toxicities, the potency to toxicity ratios are approximately equal.²⁹

Drowsiness after bupivacaine injection appears with arterial plasma concentrations greater than 1.5 ug/ml, peripheral paresthesias at 2.0 ug/ml and convulsions occur near 4.0 ug/ml.⁴⁸ Convulsions have been reported after bupivacaine injection in labor when the venous drug concentration was 2.3 ug/ml,⁵⁴ as might be expected since peripheral venous bupivacaine concentration is less than concomitant arterial plasma concentration.^{29,33} Psychomotor impairment in automobile driving appears after 1.3 ug/kg of bupivacaine is administered intramuscularly.⁵⁵

The amygdala is believed to be the primary locus of activity in local anesthetic seizures.^{29,56} Scott⁵² however, found no localized EEG changes during clinical CNS reactions to bupivacaine. Seizure threshold experiments in monkeys failed to localize aberrant EEG activity.²⁹ Short acting barbiturates and benzodiazepine derivatives are the anticonvulsants of choice for treatment of these seizures.^{29,53,56}

Hypotension due to epidural blocks occurs occasionally^{57,58} and is secondary to sympathetic blockade³⁸ or maternal caval occlusion.⁵⁸ Fetal bradycardia is a rare complication^{57,58} but Apgar scores for infants delivered with bupivacaine epidural anesthesia are not affected.⁴⁸

Previously, maximum doses of bupivacaine have been stated as 150-225 mg. every three hours and 400 mg. every 24 hours. These now appear conservative,⁵⁴ although insufficient data exists to accurately define maximum toxic doses.

Neuro toxicity with 0.5% bupivacaine has been suspected when blocks in excess of 60 hours have been achieved.⁵⁹ Another indication of local toxicity is the reduced vasodilatory response to ischemia in the presence of bupivacaine.²²

Use in Oral Surgery

Attempts to use the increased duration of action of bupivacaine to modify post operative oral surgical pain have been made.^{28,34,35,60,61} Rapid onset, profound surgical anesthesia, lack of toxic reactions, and increased duration of action have been realized. With the exception of Hellden and Associates,²⁸ studies have shown a significant delay in initial request for post operative analgesia.^{34,35,60,61} Laskin and Associates,³⁵ describe a protracted period of post operative analgesia persisting after normal sensation has returned. Local anesthetic agents in the perioral area are subject to a decreased duration of action secondary to the vascularity of the area.^{25,28} Nevertheless, obtundation of post operative oral surgical pain for up to twelve hours is reported.³⁵

Feldman and Norderam's⁶⁰ original oral surgical statistics indicated no difference in duration of action between 0.25% and 0.5% bupivacaine. More recent oral surgical studies indicate that onset, duration, and

degree of surgical anesthesia are enhanced by increasing drug concentrations.^{28,34,35,61}

Studies comparing bupivacaine with and without vasoconstrictors demonstrate no significant differences in duration between these groups.^{6,26,39,40} Prolongation of bupivacaine block in oral surgery may be achieved by combination with low molecular weight dextran -40. In one study, the mean duration of post operative analgesia in the bupivacaine-dextran group was 36 hours compared to 12 hours for the bupivacaine-saline control.⁶² The mechanism suggested is the formation of a dextran-bupivacaine complex which is absorbed much more slowly than bupivacaine alone.

Discussion

Bupivacaine is a long acting local anesthetic that is becoming extensively used in clinical practice. The potency to toxicity ratio (anesthetic index) is favorable when compared to other currently used local anesthetics.

In oral surgery, this drug provides excellent surgical anesthesia and extended post operative analgesia. Bupivacaine is ideally suited for use in providing relief of severe dental pain when definitive treatment is to be postponed. Additionally, periodontal surgery as well as prolong dental restorative procedures would benefit from bupivacaine anesthesia.

The possibilities for self-mutilation must always be considered when contemplating the use of a long-acting perioral local anesthetic.

The excellent results with bupivacaine underscore its clinical efficacy and motivate continued use and evaluation in oral surgery.

REFERENCES

1. Ekenstam B Enger B and Petterson G N-alkyl pyrrolidine and N-alkyl piperidine carboxylic acid amides Acta Chem Scand 11:183, 1957.
2. de Jong R H *Physiology and Pharmacology of Local Anesthesia* ed. 2 Springfield Illinois 1974 Charles C Thomas.
3. Moore D C Brindenbaugh L D Brindenbaugh P O and Tucker G T Bupivacaine, a review of 2,077 cases J Am Med Ass 214:713, 1960.
4. Belfrage P Raabe N Thalme B Berlin A Lumbar epidural analgesia with bupivacaine in labor. Determination of drug concentration and pH in fetal scalp blood, and continuous fetal heart rate monitoring Am J Obstet Gynecol 121 (3):360-5, 1975.
5. Moir D D Slater P J Torburn J McLaren R Moodie J Extradural analgesia in obstetrics: a controlled trial of carbonated lignocaine and bupivacaine hydrochloride with or without adrenaline Br J Anaesth 48 (2):129-35, 1976.
6. Abdel-Salam A Scott B Bupivacaine and etidocaine in epidural block for post operative relief of pain Acta Anaesthes Scand Suppl (60):08-2, 1975.
7. Phillips G A double blind trial of bupivacaine and etidocaine in epidural block for surgical induction of labour Acta Anaesthes Scand Suppl (60):94-6, 1976.
8. Kuah K B Caudal block analgesia in labouring women Med J Malaysia 28 (3):187-90, 1974.

9. Moore D C Bridenbaugh P O Bridenbaugh L D Thompson G E Balfour R I Lysons D F Bupivacaine compared with etidocaine for vaginal delivery *Anesth Analg* 54 (2):250-6, 1975.
10. Belfrage P Berlin A Lindstedt M Raabe N Plasma levels of bupivacaine following pudendal block in labour *Br J Anaesth* 45 (10):1067-9, 1973.
11. Baskett T F Carson R M Paracervical block with bupivacaine *Can Med Assoc J* 110 (12):1363-5, 1974.
12. Galindo A Hernandez J Benavides O Ortegón D E Muñoz S Bonica J J Quality of spinal extradural anaesthesia: the influence of spinal nerve root diameter *Br J Anaesth* 47 (1):41-7, 1975.
13. Moore D C Bridenbaugh P O Bridenbaugh L D Thompson G E Balfour R I Lysons D F A double-blind study of bupivacaine and etidocaine for epidural (peridural) block *Anesth Analg* 53 (5):690-7, 1974.
14. Renck H Edström H Kinnberger B Brandt G Thoracic epidural analgesia — II: Prolongation in the early post operative period by continuous injection of 1.0% bupivacaine *Acta Anaesthes Scand* 20 (1):47-56, 1976.
15. Galway J E Caves P K Dundee J W Effect of intercostal nerve blockade during operation on lung function and the relief of pain following thoracotomy *Br J Anaesth* 47 (6):730-5, 1975.
16. Willdeck-Lund G Edström H Etidocaine in intercostal nerve block for pain relief after thoracotomy A comparison with bupivacaine, *Acta Anaesthes Scand Suppl* (60):33-8, 1975.
17. Hollmen A The placental transfer of bupivacaine and lidocaine and their myocardial and cerebral uptake in the rabbit: *Acta Anaesthes Scand* 17 (3):149-52, 1973.
18. Smith D L The evaluation of chlorprocaine in combination with bupivacaine for brachial plexus anesthesia and application of nerve blockade monitor placement *J Am Osteopath Assoc* 75 (8):729-30, 1976.
19. Cunningham N L Kaplan J A A rapid onset long-acting regional anesthetic technique *Anesthes* 41 (5):509-11, 1974.
20. Radtke H Nolte H Fruhstorfer H Zenz M A comparative study between etidocaine and bupivacaine in ulnar nerve block *Acta Anaesthes Scand Suppl* (60):17-20, 1975.
21. Ware R J Intravenous regional analgesia using bupivacaine *Anaesthesia* 30 (6):817-22, 1975.
22. Drummond G B Effects of regional intravenous anesthesia on blood flow, *Br J Anaesth* 47 Suppl: 237-40, 1975.
23. Defalque R J Treatment of chronic pain with bupivacaine epidural blocks *Anesth Analg* 53 (6):841-3, 1974.
24. Adler P The use of bupivacaine for blocking the gasserian ganglion in major trigeminal neuralgia *Int J Oral Surg* 4 (6):251-7, 1975.
25. de Jong R H *Physiology and pharmacology of local anesthesia* ed 2, Springfield Illinois 1974 Charles C. Thomas.
26. Löfström B Clinical evaluation of local anesthetics *Clin Anesth* 2:19-43, 1969.
27. Henn F Brattsand R Some pharmacological and toxicological properties of a new long-acting local analgesic. LAC-43 (Marcaine), in comparison with mepivacaine and tetracaine *Acta Anaesthes Scand Suppl* 21:9, 1966.
28. Hellden L Blomberg S Woxbert B Ohman A A controlled trial of a long-acting local anesthetic (Marcaine) in oral surgery for relief of post-operative pain *Scan Tandlak Tidskr* 67 (4):223-8, 1974.
29. Munson E S Tucker W K Ausinsch B Malgodi M H Etidocaine bupivacaine and lidocaine seizure thresholds in monkeys 42 (4), 1975.
30. Renk H Edström H Thoracic epidural analgesia — a double blind study between etidocaine and bupivacaine *Acta Anaesthes Scand Suppl* (60):68-71, 1975.
31. Tammisto T Rosenberg P Tigerstedt F Comparison between bupivacaine and etidocaine in lumbar epidural block for urological surgery *Acta Anaesthes Scand Suppl* (60):68-71, 1975.
32. Niesel H C Munch I Experience with etidocaine and bupivacaine in epidural analgesia *Acta Anaesthes Scand Suppl* (60):60-3, 1975.
33. Moore D C Mather L E Bridenbaugh P O Bridenbaugh L D Balfour R I Lysons D F Horton W G Arterial and venous plasma levels of bupivacaine following epidural and intercostal nerve block *Anesthes* 45 (1):39-45, 1976.
34. Nespeca J A Clinical trials with bupivacaine in oral surgery *Oral Surg.* 42. (3):301-7, 1974.
35. Laskin J L Wallace W R DeLeo B Use of bupivacaine hydrochloride in oral surgery — a clinical study *J Oral Surg* 35, 1977.
36. Phillips G A double blind trial of bupivacaine (Marcaine) and etidocaine (Duranest) in extradural block for surgical induction of labour *Br J Anaesth* 47 (12):1305-8, 1975.
37. Collier C B Letter: Horner's Syndrome following obstetric extradural block analgesia *Br J Anaesth* 47 (2):1342, 1975.
38. Griffiths D F Diamond A W Cameron J D Postoperative extradural analgesia following thoracic surgery A Feasibility Study *Br J Anaesth* 47 (1):48-55, 1975.
39. Wilson J A double-blind comparison of single doses of etidocaine 1% and bupivacaine 0.5% during continuous lumbar epidural block in obstetrics *Acta Anaesthes Scand Suppl* (60):97-9, 1975.
40. Gitschmann J Nolte H Comparative study with etidocaine and bupivacaine in epidural block *Acta Anaesthes Scand Suppl* (60):55-9, 1975.
41. Kier L Continuous epidural analgesia in prostatectomy: comparison of bupivacaine with and without adrenaline *Acta Anaesthes Scand* 18(1):1-4, 1974.
42. Widman B Plasma concentration of local anesthetic agents in regard to absorption, distribution, and elimination, with special reference to Bupivacaine *Br J Anaesth* 47 Suppl 231-6, 1975.
43. Mather L E Long G J Thomas J The intravenous toxicity and clearance of bupivacaine in man *Clin Pharmacol Ther* 2:935, 1971.
44. de Jong R H *Physiology and Pharmacology of Local Anesthesia*, ed. 2, Springfield Illinois 1974 Charles C. Thomas.
45. Moore D C Bridenbaugh L D Bridenbaugh P O et al Caudal and epidural blocks with bupivacaine for childbirth: Report of 657 parturients *Obstet Gynecol* 37:667, 1971.
46. Scott D B Jebson P J Boyes R N Pharmacokinetic study of the local anesthetics bupivacaine (Marcaine) and etidocaine (Duranest) in Man *Br J Anaesth* 45 (10):1010-2, 1973.
47. Sjostrand U Widman B Distribution of bupivacaine in the rabbit under normal and acidotic conditions *Acta Anaesthes Scand Suppl* 50:1-24, 1973.
48. Belfrage P Berlin A Raabe N Thalme B Lumbar epidural analgesia with bupivacaine in labor drug concentration in maternal and neonatal blood at birth and during the first day of life *Am J Obstet Gynecol* 123. (8):839-44, 1975.
49. Engberg G Holmdahl M H Edström H H A comparison of the local anesthetic properties of bupivacaine and two new long-acting agents HS 37 and etidocaine in epidural analgesia *Acta Anaesthes Scand* 18 (4):277-89, 1974.
50. Reynolds F Hargrove R L Wyman J B Maternal and foetal plasma concentrations of bupivacaine after epidural block *Br J Anaesth* 45 (10):1049-53, 1973.
51. Tucker G T Mather L E Letter: Plasma protein binding of bupivacaine and its interaction with other drugs in man *Br J Anaesth* 47 (9):1029-30, 1975.
52. Scott D B Evaluation of the toxicity of local anaesthetic agents in man *Br J Anaesth* 47 (1):56-61, 1975.
53. de Jong, R. H.: *Physiology and Pharmacology of Local Anesthesia* ed. 2, Springfield Illinois 1974 Charles C. Thomas.
54. Tyack A J Parsons R J Millar D R Nicholas A D Uterine activity and plasma bupivacaine levels after caudal epidural analgesia *J Obstet Gynaecol Br Common W* 80 (10):896-901, 1973.
55. Kortilla K Hakkinen S Linnoila M Side effects and skills related to driving after intramuscular administration of bupivacaine and etidocaine *Acta Anaesthes Scand* 19 (5):384-91, 1975.

56. de Jong, R. H.: *Physiology and Pharmacology of Local Anesthesia*, ed. 2, Springfield, Illinois, 1974, Charles C. Thomas, Chapter 11.
57. Diesbecq W Rolly G Thierry M Continuous lumbar epidural analgesia during labor and delivery *Acta Anaesthes Belg* 25 (1):22-7, 1974.
58. Maltau J M The frequency of fetal bradycardia during selective epidural anaesthesia *Acta Obstet Gynecol Scand* 54 (4):357-61, 1975.
59. Pathy G V Rosen M Prolonged block with recovery after extradural analgesia for Labour *Br J Anaesth* 47 (4):520-2, 1975.
60. Feldman C Morderam A Marcaine® in oral surgery. A clinical comparative study with carbocaine® *Acta Anaesthes Scand Suppl XXIII*:409, 1966.
61. Pricco D F An evaluation of bupivacaine for regional nerve block in oral surgery *J Oral Surg* 35, 126-129, 1977.
62. Kaplan J A Miller E R Jr Gallagher E G Jr Post operative analgesia for thoracotomy patients *Anesthe Analg* 54(6):773-7, 1975.

Anesthesia Progress subscriber service

Moving?—If you're changing your address, let us know three weeks in advance, indicating your new and old location below.

NEW ADDRESS

Name _____

Address _____

City/State/ZIP _____

OLD ADDRESS OR AFFIX OLD MAILER LABEL

Name _____

Address _____

City/State/ZIP _____

**Mail to: Secretary, Dr. Peter H. Jacobsohn
P.O. Box 2297, Iowa City
Iowa 52240**

Looking for
practical advice
to better handle
everyday clinical
problems?

Look no further:

A New Book! HANDBOOK OF MEDICAL EMERGENCIES IN THE DENTAL OFFICE. By Stanley F. Malamed, D.D.S. Rely on this new book for on-the-spot assistance in life-threatening emergencies that may occur in the dental office. Six major emergency situations are examined: unconsciousness, respiratory difficulty, local anesthetic reactions, convulsive seizures, drug reactions, and chest pains. Throughout, emphasis is on prevention. April, 1978. Approx. 416 pp., 142 illus. **About \$14.50.**

New 6th Edition! MONHEIM'S LOCAL ANESTHESIA AND PAIN CONTROL IN DENTAL PRACTICE. By C. Richard Bennett, D.D.S., Ph.D.; with 3 contributors. This new edition alerts you to all the latest developments in regional anesthesia. Discussions thoroughly examine theories of pain, anatomical considerations, techniques, dosages, complications, sterilization, armamentarium, and office emergencies. More than 70 new illustrations have been added to offer you precise depictions of procedures and techniques. January, 1978. 368 pp., 123 illus. **Price, \$17.50.**

New 2nd Edition! CONSCIOUS SEDATION IN DENTAL PRACTICE. By C. Richard Bennett, D.D.S., Ph.D.; with 2 contributors. Refresh your knowledge of various pain control methods with this comprehensive source. The author first explores pain in general, then discusses assessment of the medical and dental history, including an important section on anxiety. Subsequent chapters examine such modes for inducing conscious sedation as inhalation, oral, subcutaneous, intramuscular, intravenous, and rectal. Each chapter includes material on both theory and technique, along with well-illustrated, step-by-step descriptions. July, 1978. Approx. 256 pp., 102 illus. **About \$18.50.**

To order your on-approval copies, simply call our toll-free number: (800) 325-4177 and ask for extension 10, 9 am to 5 pm (CDT) Monday through Friday. A80451

MOSBY
TIMES MIRROR

THE C. V. MOSBY COMPANY
11830 WESTLINE INDUSTRIAL DRIVE
ST. LOUIS, MISSOURI 63141